

Critical points of the N -vortex Hamiltonian in bounded planar domains and steady state solutions of the incompressible Euler equations

THOMAS BARTSCH*[†]

ANGELA PISTOIA[‡]

Abstract

We prove the existence of critical points of the N -vortex Hamiltonian

$$H_{KR}(x_1, \dots, x_N) = \sum_{i=1}^N \Gamma_i^2 h(x_i) + \sum_{\substack{i,j=1 \\ j \neq i}}^N \Gamma_i \Gamma_j G(x_i, x_j) + 2 \sum_{i=1}^N \Gamma_i \psi_0(x_i)$$

in a bounded domain $\Omega \subset \mathbb{R}^2$ which may be simply or multiply connected. Here G denotes the Green function for the Dirichlet Laplace operator in Ω , more generally a hydrodynamic Green function, and h the Robin function. Moreover $\psi_0 \in C^1(\overline{\Omega})$ is a harmonic function on Ω . The domain need not be simply connected. We obtain new critical points $x = (x_1, \dots, x_N)$ for $N = 3$ or $N = 4$ under conditions on the vorticities $\Gamma_i \in \mathbb{R} \setminus \{0\}$. These critical points correspond to point vortex equilibria of the Euler equation in vorticity form. The case $\Gamma_i = (-1)^i$ of counter-rotating vortices with identical vortex strength is included. The point vortex equilibria can be desingularized to obtain smooth steady state solutions of the Euler equations for an ideal fluid. The velocity of these steady states will be irrotational except for N vorticity blobs near x_1, \dots, x_N .

Keywords: vortex dynamics, point vortices, counter-rotating vortices, steady states of the Euler flow

AMS subject classification: 35J60, 35J25, 37J45, 76B47.

1 Introduction

The dynamics of N point-vortices $x_1, \dots, x_N \in \Omega$ in a bounded domain $\Omega \subset \mathbb{R}^2$ in the plane is governed by a Hamiltonian system

$$(1.1) \quad \begin{cases} \Gamma_i \frac{dx_{i1}}{dt} = \frac{\partial H_{KR}}{\partial x_{i2}}(x_1, \dots, x_N); \\ \Gamma_i \frac{dx_{i2}}{dt} = -\frac{\partial H_{KR}}{\partial x_{i1}}(x_1, \dots, x_N); \end{cases} \quad i = 1, \dots, N.$$

Here $\Gamma_i \in \mathbb{R} \setminus \{0\}$ denotes the strength of the i -th vortex x_i , the sign determining the orientation of the vortex. The Hamiltonian is given by the Kirchhoff-Routh path function

$$(1.2) \quad H_{KR}(x_1, \dots, x_N) = \sum_{i=1}^N \Gamma_i^2 h(x_i) + \sum_{\substack{i,j=1 \\ i \neq j}}^N \Gamma_i \Gamma_j G(x_i, x_j) + 2 \sum_{i=1}^N \Gamma_i \psi_0(x_i)$$

*T.B. thanks the Università di Roma “La Sapienza” for the invitation and the hospitality during several visits.

[†]Supported by DAAD grant 50766047.

[‡]Supported by the M.I.U.R. National Project “Metodi variazionali e topologici nello studio di fenomeni non lineari”.

where

$$G(x, y) = g(x, y) - \frac{1}{2\pi} \log |x - y|$$

is the Green function of the Dirichlet Laplacian in Ω . Here $g : \Omega \times \Omega \rightarrow \mathbb{R}$ is the regular part, and $h : \Omega \rightarrow \mathbb{R}$, $h(x) = g(x, x)$, denotes the Robin function. Moreover $\psi_0 \in \mathcal{C}^1(\overline{\Omega})$ is a harmonic function on Ω modeling the boundary flux. In case of a solid boundary one has $\psi_0 = 0$. H_{KR} is defined on the configuration space

$$\mathcal{F}_N \Omega = \{(x_1, \dots, x_N) \in \Omega^N : x_i \neq x_j \text{ for } i \neq j\}.$$

The domain Ω need neither be simply connected nor symmetric. More generally, G can be a hydrodynamic Green function (see [13]), or even a function having certain properties of Green functions.

Based on first ideas of Helmholtz [14] about vortices, the system has been deduced by Kirchhoff [15], Routh [23], and Lin [18, 19] from the Euler equations

$$(1.3) \quad \begin{cases} v_t + (v \cdot \nabla)v = -\nabla P \\ \nabla \cdot v = 0 \end{cases}$$

for an incompressible and non-viscous fluid in Ω . Here v denotes the velocity field and P the pressure of the fluid. The scalar vorticity $\omega = \nabla \times v = \partial_1 v_2 - \partial_2 v_1$ satisfies the equation

$$(1.4) \quad \omega_t + v \cdot \nabla \omega = 0.$$

The point vortex ansatz $\omega = \sum_{k=1}^N \Gamma_k \delta_{x_k}$, where δ_{x_k} is the usual Dirac delta, leads to (1.1) for the point vortices $x_k(t)$. We refer to [13, 20–22, 24] for modern treatments of vorticity methods.

There are many results about point vortex dynamics if $\Omega = \mathbb{R}^2$ is the plane, or if Ω is a special domain like the disc, the half-disc, an annulus, an infinite strip. In these cases the Green function, hence the Hamiltonian, is either explicitly known or one has good representations of it. There are also many results of numerical nature, due to the multiple applications of point vortex methods in science and engineering. We just refer to the surveys [1, 2, 22] and the literature cited therein.

In this paper we present new conditions on the vortex strengths Γ_i such that H_{KR} has a critical point. Our results extend considerably earlier ones from [4, 6, 10] where only special cases have been treated, all dealing with $\Gamma_i \in \{\pm 1\}$ and $\psi_0 = 0$. Observe that $\mathcal{F}_N \Omega \subset \Omega^N$ is an open bounded subset of \mathbb{R}^{2N} , and that H_{KR} is singular and not bounded from above nor below. Therefore the existence of critical points is highly nontrivial, in particular since we require no symmetry nor any geometrical or topological properties of the domain. Our results hold for functions $F : \mathcal{F}_N \Omega \rightarrow \mathbb{R}$ which are \mathcal{C}^1 -close to H_{KR} on certain compact subsets of $\mathcal{F}_N \Omega$. This allows to apply the methods from Cao, Liu and Wei [7, 8] on the desingularization of stationary point vortex solutions and to obtain stationary solutions of the Euler equations (1.3), (1.4). This is done by constructing families ψ_ε of stream functions with vortex blobs which converge as $\varepsilon \rightarrow 0$ towards the stationary point vortices we construct. The velocity v will be irrotational outside these vortex blobs.

The paper is organized as follows. First, in Section 2 we state our main results Theorems 2.1 to 2.3 about the existence of critical points of *Hamiltonians of the N -vortex type*, and we state in Theorem 2.4 our results about solutions of the incompressible Euler equations. Next, in Section 3 we prove a compactness result for the class of Hamiltonians we consider. This is very technical but in a sense the core of our paper. Section 4 contains the proofs of Theorems 2.1 to 2.3. Finally in Section 5 we desingularize the stationary point vortex solutions by proving Theorem 2.4.

2 Statement of results

Let $\Omega \subset \mathbb{R}^2$ be a bounded domain with \mathcal{C}^2 -boundary. We fix $\varepsilon_0 > 0$ small so that the reflection at $\partial\Omega$ is well defined in $\Omega_0 := \{x \in \Omega : \text{dist}(x, \partial\Omega) < \varepsilon_0\}$ and maps to the complement of Ω ; we denote it by

$\Omega_0 \rightarrow \mathbb{R}^2 \setminus \overline{\Omega}$, $x \mapsto \bar{x}$. It is of class \mathcal{C}^1 since $\partial\Omega$ is of class \mathcal{C}^2 . We write

$$p : \Omega_0 \rightarrow \partial\Omega, \quad p(x) = \frac{1}{2}(x + \bar{x}),$$

for the orthogonal projection onto the boundary, and

$$\nu : \Omega_0 \rightarrow \mathbb{R}^2, \quad \nu(x) = \frac{1}{|x - \bar{x}|}(x - \bar{x}),$$

for the interior normal; more precisely, $\nu(x)$ is the interior unit normal at $p(x) \in \partial\Omega$ for $x \in \Omega_0$. Clearly, $p(x) = x - \text{dist}(x, \partial\Omega)\nu(x)$ and $\bar{x} = x - 2\text{dist}(x, \partial\Omega)\nu(x)$.

Let $N \geq 2$ and $\Gamma_1, \dots, \Gamma_N \in \mathbb{R} \setminus \{0\}$ be given. We consider a *Hamiltonian of the N -vortex type*, i. e. a function $H : \mathcal{F}_N\Omega \rightarrow \mathbb{R}$ of the form

$$(2.1) \quad H(x) = \sum_{i=1}^N \Gamma_i^2 h(x_i) + \sum_{\substack{i,j=1 \\ j \neq k}}^N \Gamma_i \Gamma_j G(x_i, x_j) + f(x)$$

where $f \in \mathcal{C}^1(\overline{\Omega}^N)$ and

$$(2.2) \quad G(x, y) = g(x, y) - \frac{1}{2\pi} \log |x - y|$$

is a *generalized Green's function* by which we mean that the following properties hold.

- (A1) G is bounded from below and symmetric, i. e. $G(x, y) = G(y, x)$.
- (A2) $g : \Omega \times \Omega \rightarrow \mathbb{R}$ is a \mathcal{C}^1 -function, bounded from above, and $h(x) = g(x, x) \rightarrow -\infty$ as $\text{dist}(x, \partial\Omega) \rightarrow 0$.
- (A3) For every $\varepsilon > 0$ there is a constant $C_1 = C_1(\Omega, \varepsilon) > 0$ such that

$$|h(x)| + |\nabla h(x)| \leq C_1 \quad \text{for every } x \in \Omega \text{ with } \text{dist}(x, \partial\Omega) \geq \varepsilon$$

and

$$|G(x, y)| + |\nabla_x G(x, y)| + |\nabla_y G(x, y)| \leq C_1 \quad \text{for every } x, y \in \Omega \text{ with } |x - y| \geq \varepsilon.$$

- (A4) There exists a constant $C_2 = C_2(\Omega) > 0$ such that $\psi(x, y) := g(x, y) - \frac{1}{2\pi} \log |\bar{x} - y|$ satisfies

$$|\psi(x, y)| + |\nabla_x \psi(x, y)| + |\nabla_y \psi(x, y)| \leq C_2 \quad \text{for every } x, y \in \Omega_0.$$

It is well known that these assumptions hold for the Dirichlet Green's function, more generally for a hydrodynamic Green's function (see [13] for the definition); details can be found in [6, 16]. Our first theorem deals with a rather simple case.

Theorem 2.1. *Suppose $N = 2$ and $\Gamma_1 \Gamma_2 < 0$. There exists a compact subset $K \subset \mathcal{F}_2(\Omega)$ and $\delta > 0$ such that the following holds:*

- a) Any \mathcal{C}^1 -function $F : \mathcal{F}_2(\Omega) \rightarrow \mathbb{R}$ with $\|F|_K - H|_K\|_\infty < \delta$ has at least $\text{cat}(\mathcal{F}_2(\Omega))$ critical points (x_1^i, x_2^i) , $i = 1, \dots, \text{cat}(\mathcal{F}_2(\Omega))$ in K .
- b) If $\Gamma_1 = -\Gamma_2$ and if F is symmetric, i. e. $F(x, y) = F(y, x)$, then F has at least $k := \text{cat}(\mathcal{F}_2(\Omega)/(x_1, x_2) \sim (x_2, x_1))$ pairs $(x_1^i, x_2^i), (x_2^i, x_1^i)$ of critical points in K , $i = 1, \dots, k$.

c) If $F_\varepsilon : \mathcal{F}_2(\Omega) \rightarrow \mathbb{R}$ is a family of \mathcal{C}^1 -functions such that $\|F_\varepsilon|_K - H|_K\|_{\mathcal{C}^1} \rightarrow 0$ then the critical points x_ε obtained in a) or b) converge along a subsequence towards a critical point of H .

Here cat denotes the Lusternik-Schnirelman category. The problem becomes considerably more difficult if $N > 2$. We only deal with the cases $N = 3$, $N = 4$ and require the following assumption:

$$(2.3) \quad \begin{aligned} & \Gamma_i \Gamma_{i+1} < 0 \text{ for } i = 1, \dots, N-1, \text{ and} \\ & \text{for every subset } I \subset \{1, \dots, N\} \text{ with } |I| \geq 3 \text{ there holds } \sum_{i,j \in I, i \neq j} \Gamma_i \Gamma_j < 0. \end{aligned}$$

Theorem 2.2. *Let $N = 3$ and assume (2.3). Then there exists a compact subset $K \subset \mathcal{F}_3(\Omega)$ and $\delta > 0$ such that the following holds:*

- a) Any \mathcal{C}^1 -function $F : \mathcal{F}_3(\Omega) \rightarrow \mathbb{R}$ with $\|F|_K - H|_K\|_{\mathcal{C}^1} < \delta$ has a critical point in K .
- b) If $F_\varepsilon : \mathcal{F}_2(\Omega) \rightarrow \mathbb{R}$ is a family of \mathcal{C}^1 -functions such that $\|F_\varepsilon|_K - H|_K\|_{\mathcal{C}^1} \rightarrow 0$ then the critical points x_ε obtained in a) converge along a subsequence towards a critical point of H .

In the case $N = 4$ we need an additional hypothesis on the vorticities:

$$(2.4) \quad |\Gamma_2| < |\Gamma_1| + |\Gamma_3| \quad \text{and} \quad |\Gamma_3| < |\Gamma_2| + |\Gamma_4|.$$

Theorem 2.3. *Let $N = 4$ and assume (2.3), (2.4). Then there exists a compact subset $K \subset \mathcal{F}_4(\Omega)$ and $\delta > 0$ such that the following holds:*

- a) Any \mathcal{C}^1 -function $F : \mathcal{F}_4(\Omega) \rightarrow \mathbb{R}$ with $\|F|_K - H|_K\|_{\mathcal{C}^1} < \delta$ has a critical point in K .
- b) If $F_\varepsilon : \mathcal{F}_2(\Omega) \rightarrow \mathbb{R}$ is a family of \mathcal{C}^1 -functions such that $\|F_\varepsilon|_K - H|_K\|_{\mathcal{C}^1} \rightarrow 0$ then the critical points x_ε obtained in a) converge along a subsequence towards a critical point of H .

Observe that (2.3) and (2.4) hold if $\Gamma_i = (-1)^i$. This case has already been treated in [6]. The proof of [6, Theorem 1.2] has a gap, however, which is being fixed in this paper using a different method though. Related results concerning point vortex equilibria on general bounded domains can also be found in [16] and, if the domain is symmetric, in [17]. These papers complement our results in that different conditions on the set of vorticities are considered. Earlier results dealing with the case of Ω not being simply connected and all $\Gamma_i = 1$ can be found in [9, 12]. Periodic solutions of (HS) for any given N with all $\Gamma_i = 1$, on bounded and unbounded domains, have been constructed in [5].

The point vortex equilibria obtained in Theorems 2.1-2.3 can be regularized as limits of vorticity distributions of smooth steady state solutions of the incompressible Euler equations in the following way. Let G be the Green function of $-\Delta$ in Ω with homogeneous Dirichlet boundary conditions and let $\psi_0 \in \mathcal{C}^2(\overline{\Omega})$ be harmonic in Ω . We consider the Kirchhoff-Routh path function $H_{KR} : \mathcal{F}_N \Omega \rightarrow \mathbb{R}$ defined by

$$(2.5) \quad H_{KR}(x) = \sum_{i=1}^N \Gamma_i^2 h(x_i) + \sum_{\substack{i,j=1 \\ i \neq j}}^N \Gamma_i \Gamma_j G(x_i, x_j) + 2 \sum_{i=1}^N \Gamma_i \psi_0(x_i).$$

We write $\frac{\partial \psi_0}{\partial \tau} : \partial\Omega \rightarrow \mathbb{R}^2$ for the tangential derivative of ψ_0 on $\partial\Omega$, and we set $(w_1, w_2)^\perp = J(w_1, w_2) := (w_2, -w_1)$.

Theorem 2.4. *Consider one of the cases*

- (i) $N = 2$ and $\Gamma_1 \Gamma_2 < 0$;

(ii) $N = 3$ and (2.3) holds;

(iii) $N = 4$ and (2.3), (2.4) hold.

Then for $\varepsilon > 0$ small there exists a stationary solution $v_\varepsilon : \Omega \rightarrow \mathbb{R}^2$ of (1.3) with pressure P_ε and boundary flux $v(x) \cdot \nu(x) = \frac{\partial \psi_0(x)}{\partial \tau}$. Moreover, the scalar vorticity of v_ε is of the form $\omega_\varepsilon = \nabla \times v_\varepsilon = \sum_{i=1}^N \omega_{i,\varepsilon}$ with $\text{supp}(\omega_{i,\varepsilon}) \rightarrow x_i^* \in \Omega$ as $\varepsilon \rightarrow 0$ along a subsequence, $\int_\Omega \omega_{i,\varepsilon} \rightarrow \Gamma_i$, where $(x_1^*, \dots, x_N^*) \in \mathcal{F}_N \Omega$ is a critical point of the Kirchhoff-Routh path function H_{KR} from (2.5).

Here $\text{supp}(\omega_{i,\varepsilon}) \rightarrow x_i \in \Omega$ means that for $\delta > 0$ the support $\text{supp}(\omega_{i,\varepsilon})$ is contained in the δ -neighborhood of $x_i \in \Omega$ provided ε is small. Theorem 2.4 will be proved by the method of stream functions. Recall that a stream function $\psi : \Omega \rightarrow \mathbb{R}$ for v satisfies $v = J\nabla\psi = (-\partial\psi/\partial x_2, \partial\psi/\partial x_1)$, hence $\omega = -\Delta\psi$ and $v = J\nabla(-\Delta)^{-1}\omega$. If $\psi : \Omega \rightarrow \mathbb{R}$ satisfies

$$(2.6) \quad \begin{cases} -\Delta\psi = F'(\psi) & \text{for } x \in \Omega, \\ \psi = \psi_0 & \text{for } x \in \partial\Omega, \end{cases}$$

for some arbitrary function $F \in C^2(\mathbb{R})$ then $v = J\nabla\psi$ solves (1.3) with pressure field $P = F(\psi) - \frac{1}{2}|\nabla\psi|^2$ and vorticity $F'(\psi)$. Using the method from [7, 8] and our Theorems (2.1)–(2.3) there are appropriate functions F_ε and solutions of (2.6) with $F = F_\varepsilon$ which will yield Theorem 2.4. The theorems from [7, 8] cannot be applied directly because there it is assumed that the Kirchhoff-Routh path function H_{KR} has an isolated stable critical point. This will not be the case in general, for instance, it doesn't hold for Ω a disc or an annulus. The latter case is excluded in [7] anyway because there the domain is required to be simply connected. This is needed when one wants to prescribe the boundary flux, not the function ψ_0 .

3 A compactness result

We fix a function G as in (2.2) such that (A1)–(A4) hold, we fix a function $f \in C^1(\overline{\Omega})$, and we consider a Hamiltonian H as in (2.1). Then we introduce the function $\Phi : \mathcal{F}_N \Omega \rightarrow \mathbb{R}$ defined by

$$\Phi(x) := \sum_{i=1}^N \Gamma_i^2 h(x_i) - \sum_{\substack{i,j=1 \\ i \neq j}}^N |\Gamma_i \Gamma_j| G(x_i, x_j).$$

Assumptions (A1) and (A2) imply

$$\lim_{x \rightarrow \partial \mathcal{F}_N \Omega} \Phi(x) = -\infty.$$

Proposition 3.1. *Assume that $N \in \{3, 4\}$ and (2.3) is satisfied. Then for any $a, b \in \mathbb{R}$ with $a < b$ there exists $M_0 > 0$ such that the following holds:*

$$\Phi(x) \leq -M_0, \ a \leq H(x) \leq b, \ \nabla H(x) = \lambda \nabla \Phi(x) \quad \implies \quad \lambda > 0.$$

The rest of this section is concerned with the proof of Proposition 3.1. We argue by contradiction. Suppose there exist $a, b \in \mathbb{R}$ with $a < b$, a sequence of points $x^n = (x_1^n, \dots, x_N^n) \in \mathcal{F}_N \Omega$, and a sequence $\lambda_n \leq 0$ such that

$$(3.1) \quad \Phi(x^n) \rightarrow -\infty, \ a \leq H(x^n) \leq b, \ \text{and } \nabla H(x^n) = \lambda_n \nabla \Phi(x^n).$$

Recall from Section 2 the reflection $x \mapsto \bar{x}$ at the boundary, the projection $x \mapsto p(x)$ onto the boundary, and the interior normal $x \mapsto \nu(x)$. These maps are defined for $x \in \Omega_0$ close to the boundary. We set $d_i^n := \text{dist}(x_i^n, \partial\Omega)$, and $\nu_i^n := \nu(x_i^n)$, $p_i^n := p(x_i^n)$, if $x_i^n \in \Omega_0$. In the sequel $O(1)$, $o(1)$ refer to $n \rightarrow \infty$. The following lemma holds for all sequences $(x^n)_n$ in $\mathcal{F}_N \Omega$.

- Lemma 3.2.** (i) $h(x_i^n) = \frac{1}{2\pi} \log 2d_i^n + O(1)$ and $d_i^n |\nabla h(x_i^n)| = O(1)$ if $x_i^n \in \Omega_0$.
(ii) $\nabla h(x_i^n) = \frac{1}{2\pi d_i^n} \nu_i^n + o(1)$ if $d_i^n \rightarrow 0$
(iii) $G(x_i^n, x_j^n) = -\frac{1}{2\pi} \log |x_i^n - x_j^n| + \frac{1}{2\pi} \log |x_i^n - \bar{x}_j^n| + O(1)$ if $x_j^n \in \Omega_0$.
(iv) $G(x_i^n, x_j^n) = O(1)$ if $\liminf \frac{|x_i^n - x_j^n|}{d_i^n} > 0$.
(v) $\partial_1 G(x_i^n, x_j^n) = -\frac{1}{2\pi} \left(\frac{x_i^n - x_j^n}{|x_i^n - x_j^n|^2} + \frac{\bar{x}_i^n - x_j^n}{|\bar{x}_i^n - x_j^n|^2} \right) + O(1) = -\frac{1}{2\pi} \left(\frac{x_i^n - x_j^n}{|x_i^n - x_j^n|^2} + \frac{x_i^n - \bar{x}_j^n}{|x_i^n - \bar{x}_j^n|^2} \right) + O(1)$ if $x_i^n \in \Omega_0$ or $x_j^n \in \Omega_0$, respectively.
(vi) $d_i^n |\nabla g(x_i^n, x_j^n)| = O(1)$ if $x_i^n \in \Omega_0$.
(vii) $\langle \partial_1 G(x_i^n, x_j^n), \nu_i^n \rangle + \langle \partial_1 G(x_j^n, x_i^n), \nu_j^n \rangle = \frac{1}{2\pi} (d_i^n + d_j^n) \left(\frac{1}{|\bar{x}_i^n - x_j^n|^2} + \frac{1}{|\bar{x}_j^n - x_i^n|^2} \right) + O(1)$ if $x_i^n, x_j^n \in \Omega_0$.
(viii) $|\bar{x}_i^n - x_j^n|^2 = |x_i^n - x_j^n|^2 + 4d_i^n d_j^n + o(|x_i^n - x_j^n|^2)$ if $x_i^n, x_j^n \rightarrow x^* \in \partial\Omega$.
(ix) $\langle p_i^n - p_j^n, \nu_i^n \rangle = O(|x_i^n - x_j^n|^2)$ if $x_i^n, x_j^n \in \Omega_0$.

Proof. These statements follow in a straightforward way from assumptions (A1)–(A4). \square

We write the proof of Proposition 3.1 for $N = 4$. The case $N = 3$ is simpler and can be deduced by forgetting all arguments which involve x_4^n . In the sequel we drop the notation $n \rightarrow \infty$ from all kinds of limits. The first lemma does not require hypothesis (2.3). It is sufficient that all $\Gamma_i \neq 0$ for $i = 1, \dots, 4$.

Lemma 3.3. *There exist indices $i_0 \neq j_0$ such that $\liminf \frac{|x_{i_0}^n - x_{j_0}^n|}{d_{i_0}^n} \rightarrow 0$.*

Proof. Suppose to the contrary that $|x_i^n - x_j^n| \geq cd_i^n$ for all $i \neq j$. Then (3.1) implies that $d_k^n \rightarrow 0$ for some $k \in \{1, \dots, 4\}$. Using Lemma 3.2 we can estimate the energy:

$$\begin{aligned} H(x^n) &= \sum_{i=1}^N \Gamma_i^2 h(x_i^n) + \sum_{\substack{i,j=1 \\ i \neq j}}^N \Gamma_i \Gamma_j G(x_i^n, x_j^n) = \sum_{i=1}^N \Gamma_i^2 h(x_i^n) + O(1) \\ &\leq \Gamma_k^2 h(x_k^n) + O(1) = \frac{1}{2\pi} \ln d_k^n + O(1) \rightarrow -\infty. \end{aligned}$$

This contradicts (3.1). \square

After passing to a subsequence we may assume for each $i \in \{1, \dots, 4\}$:

$$(3.2) \quad \text{either } |x_i^n - x_{i_0}^n| = o(d_{i_0}^n) \quad \text{or} \quad \liminf \frac{|x_i^n - x_{i_0}^n|}{d_{i_0}^n} > 0.$$

Setting

$$I := \{i \in \{1, \dots, 4\} : |x_i^n - x_{i_0}^n| = o(d_{i_0}^n)\}$$

Lemma 3.3 implies

$$(3.3) \quad \begin{cases} |I| \geq 2, \frac{d_i^n}{d_j^n} \rightarrow 1 \text{ and } |x_i^n - x_j^n| = o(d_i^n) \text{ for } i, j \in I, \\ |x_i^n - x_j^n| = o(|x_i^n - x_k^n|) \text{ for } i, j \in I, k \notin I. \end{cases}$$

Lemma 3.4. *The only possibilities for I are $\{1, 3\}$ or $\{2, 4\}$. Moreover $\lambda_n \rightarrow -1$.*

Proof. We set

$$z^n := (z_1^n, \dots, z_4^n), \quad z_i^n := \begin{cases} x_i^n - x_{i_0}^n & i \in I, \\ 0 & i \notin I, \end{cases}$$

and compute, using (A1)–(A4), Lemma 3.2, as well as (3.2) and (3.3),

$$\begin{aligned} \langle \nabla H(x^n), z^n \rangle &= \sum_{i \in I} \Gamma_i^2 \langle \nabla h(x_i^n), z_i^n \rangle + 2 \sum_{i \in I} \sum_{j \neq i} \Gamma_i \Gamma_j \langle \partial_1 g(x_i^n, x_j^n), z_i^n \rangle \\ &\quad - \frac{1}{\pi} \sum_{i \in I} \sum_{j \neq i} \Gamma_i \Gamma_j \frac{\langle x_i^n - x_j^n, z_i^n \rangle}{|x_i^n - x_j^n|^2} \\ &= -\frac{1}{\pi} \sum_{i, j \in I, i < j} \Gamma_i \Gamma_j \frac{\langle x_i^n - x_j^n, x_i^n - x_1^n \rangle}{|x_i^n - x_j^n|^2} + o(1) \\ &= -\frac{1}{\pi} \sum_{i, j \in I, i < j} \Gamma_i \Gamma_j + o(1) \end{aligned}$$

Arguing in the same way, we also obtain that

$$\langle \nabla \Phi(x^n), z^n \rangle = \frac{1}{\pi} \sum_{i, j \in I, i < j} |\Gamma_i \Gamma_j| + o(1).$$

Now the equation $\nabla H(x^n) = \lambda_n \nabla \Phi(x^n)$ implies

$$0 \geq \lambda_n \rightarrow -\frac{\sum_{i, j \in I, i < j} \Gamma_i \Gamma_j}{\sum_{i, j \in I, i < j} |\Gamma_i \Gamma_j|}.$$

This implies $\sum_{i, j \in I, i < j} \Gamma_i \Gamma_j \geq 0$, hence $|I| \leq 2$ by hypothesis (2.3). Now (3.3) yields $|I| = 2$, and since $\Gamma_i \Gamma_{i+1} < 0$ we must have $I = \{1, 3\}$ or $I = \{2, 4\}$. We also obtain immediately $\lambda_n \rightarrow -1$. \square

Lemma 3.5. *At least one of the following is true:*

- (i) $I = \{1, 3\}$ satisfies (3.3) and $d_1^n \rightarrow 0$.
- (ii) $I = \{2, 4\}$ satisfies (3.3) and $d_2^n \rightarrow 0$.

Proof. Suppose $I_1 = \{1, 3\}$ satisfies (3.3) but, after passing to a subsequence, $d_1^n \geq c > 0$. Since $2, 4 \notin I_1$ there holds $|x_i^n - x_j^n| \geq c$ for $i \in \{1, 3\}$, $j \in \{2, 4\}$. Now $H(x^n) = O(1)$ implies $h(x_2^n) \rightarrow -\infty$ or $h(x_4^n) \rightarrow -\infty$, hence $d_2^n \rightarrow 0$ or $d_4^n \rightarrow 0$. Assuming without loss of generality $d_2^n \rightarrow 0$, we consider the equation

$$\langle \partial_{x_2} (H(x^n) - \lambda_n \Phi(x^n)), \nu_2^n \rangle = 0.$$

Using $\partial_1 G(x_2^n, x_1^n) = O(1) = \partial_1 G(x_2^n, x_3^n)$ and $\lambda_n \rightarrow -1$ we deduce

$$(1 - \lambda_n) \frac{\Gamma_2^2}{2\pi d_2^n} + (1 + \lambda_n) \Gamma_2 \Gamma_4 \partial_1 G(x_2^n, x_4^n) + O(1) = 0$$

and therefore

$$\frac{\Gamma_2^2}{2\pi} + \Gamma_2 \Gamma_4 \frac{1 + \lambda_n}{1 - \lambda_n} \frac{\langle x_2^n - x_4^n, d_2^n \nu_2^n \rangle}{|x_2^n - x_4^n|^2} = o(1).$$

This implies $|x_2^n - x_4^n| = o(d_2^n)$. Then (3.3) holds for $I_2 = \{2, 4\}$, and $d_2^n \rightarrow 0$. \square

Without loss of generality we may now assume $I = \{1, 3\}$ and $d_1^n \rightarrow 0$. Thus there holds:

$$(3.4) \quad \begin{cases} \frac{d_1^n}{d_3^n} \rightarrow 1; & |x_1^n - x_3^n| = o(d_1^n) = o(d_3^n); \\ x_1^n, x_3^n \rightarrow p \in \partial\Omega; & |x_i^n - x_1^n| \geq cd_1^n \text{ for } i \in \{2, 4\}. \end{cases}$$

After passing to a subsequence we can also assume for $i \in \{1, 3\}$:

$$(3.5) \quad \frac{d_1^n}{|x_i^n - x_2^n|} \rightarrow \alpha_1; \quad \frac{d_2^n}{|x_i^n - x_2^n|} \rightarrow \alpha_2; \quad \frac{d_1^n}{|x_i^n - x_4^n|} \rightarrow \beta_1; \quad \frac{d_2^n}{|x_i^n - x_4^n|} \rightarrow \beta_2.$$

Clearly we have $\alpha_1, \alpha_2, \beta_1, \beta_2 \geq 0$.

Lemma 3.6. Fix $i \in \{1, 3\}$ and suppose $x_2^n \rightarrow p \in \partial\Omega$. Then there holds:

$$\begin{aligned} (i) \quad & d_i^n \frac{\langle x_i^n - x_2^n, \nu_i^n \rangle}{|x_i^n - x_2^n|^2} \rightarrow \alpha_1(\alpha_1 - \alpha_2) \\ (ii) \quad & d_i^n \frac{\langle x_i^n - \bar{x}_2^n, \nu_i^n \rangle}{|x_i^n - \bar{x}_2^n|^2} \rightarrow \frac{\alpha_1(\alpha_1 + \alpha_2)}{1 + 4\alpha_1\alpha_2} \\ (iii) \quad & \frac{\langle \nu_2^n, x_2^n - x_i^n \rangle}{d_2^n} \rightarrow 1 - \frac{\alpha_1}{\alpha_2} \text{ provided } \alpha_2 > 0. \\ (iv) \quad & \frac{\langle x_2^n - \bar{x}_i^n, x_2^n - x_i^n \rangle}{|x_2^n - \bar{x}_i^n|^2} \rightarrow \frac{1 + 2\alpha_1(\alpha_2 - \alpha_1)}{1 + 4\alpha_1\alpha_2} \end{aligned}$$

Proof. We compute using Lemma 3.2:

$$\begin{aligned} d_i^n \frac{\langle x_i^n - x_2^n, \nu_i^n \rangle}{|x_i^n - x_2^n|^2} &= d_i^n \frac{\langle d_i^n \nu_i^n - d_2^n \nu_2^n, \nu_i^n \rangle}{|x_i^n - x_2^n|^2} + o(1) \\ &= \frac{|d_i^n|^2}{|x_i^n - x_2^n|^2} - \frac{d_i^n d_2^n}{|x_i^n - x_2^n|^2} (1 + \langle \nu_2^n - \nu_i^n, \nu_i^n \rangle) + o(1) \\ &\rightarrow \alpha_1(\alpha_1 - \alpha_2), \end{aligned}$$

This proves (i). Next, (ii) follows from:

$$\begin{aligned} d_i^n \frac{\langle x_i^n - \bar{x}_2^n, \nu_i^n \rangle}{|x_i^n - \bar{x}_2^n|^2} &= d_i^n \frac{\langle d_i^n \nu_i^n + d_2^n \nu_2^n, \nu_i^n \rangle}{|x_i^n - \bar{x}_2^n|^2 + 4d_i^n d_2^n + o(|x_i^n - x_2^n|^2)} + o(1) \\ &\rightarrow \frac{\alpha_1(\alpha_1 + \alpha_2)}{1 + 4\alpha_1\alpha_2}. \end{aligned}$$

In order to see (iii) we calculate:

$$\begin{aligned} \frac{\langle \nu_2^n, x_2^n - x_i^n \rangle}{d_2^n} &= \frac{\langle \nu_2^n, p_2^n - p_i^n \rangle}{|x_2^n - \bar{x}_i^n|} \cdot \frac{|x_2^n - \bar{x}_i^n|}{d_2^n} + \frac{\langle \nu_2^n, d_2^n \nu_2^n - d_i^n \nu_i^n \rangle}{d_2^n} \\ &= o(1) \cdot \frac{1}{\alpha_2} + 1 - \frac{d_i^n}{d_2^n} + \frac{d_i^n \langle \nu_2^n, \nu_2^n - \nu_i^n \rangle}{d_2^n} \\ &\rightarrow 1 - \frac{\alpha_1}{\alpha_2} \end{aligned}$$

Finally we prove (iv):

$$\begin{aligned}
\frac{\langle x_2^n - \bar{x}_i^n, x_2^n - x_i^n \rangle}{|x_2^n - \bar{x}_i^n|^2} &= \frac{|x_2^n - \bar{x}_i^n|^2 + \langle 2d_i^n \nu_i^n, x_2^n - x_i^n \rangle}{|x_2^n - \bar{x}_i^n|^2 + 4d_i^n d_2^n + o(|x_2^n - x_i^n|^2)} \\
&= \frac{|x_2^n - \bar{x}_i^n|^2 + 2d_i^n \langle \nu_i^n, d_2^n \nu_2^n - d_i^n \nu_i^n \rangle + o(|x_2^n - x_i^n|^2)}{|x_2^n - \bar{x}_i^n|^2 + 4d_i^n d_2^n + o(|x_2^n - x_i^n|^2)} \\
&\rightarrow \frac{1 + 2\alpha_1(\alpha_2 - \alpha_1)}{1 + 4\alpha_1\alpha_2}.
\end{aligned}$$

□

We also need the following equality:

$$\begin{aligned}
(3.6) \quad 0 &= \langle \partial_{x_1} (H(x^n) - \lambda_n \Phi(x^n)), \nu_1^n \rangle + \langle \partial_{x_3} (H(x^n) - \lambda_n \Phi(x^n)), \nu_3^n \rangle \\
&= (1 - \lambda_n) \left(\frac{\Gamma_1^2}{2\pi d_1^n} + \frac{\Gamma_3^2}{2\pi d_3^n} - 2|\Gamma_1 \Gamma_2| \langle \partial_1 G(x_1^n, x_2^n), \nu_1^n \rangle - 2|\Gamma_1 \Gamma_4| \langle \partial_1 G(x_1^n, x_4^n), \nu_1^n \rangle \right. \\
&\quad \left. - 2|\Gamma_3 \Gamma_2| \langle \partial_1 G(x_3^n, x_2^n), \nu_3^n \rangle - 2|\Gamma_3 \Gamma_4| \langle \partial_1 G(x_3^n, x_4^n), \nu_3^n \rangle \right) \\
&\quad + 2(1 + \lambda_n) \Gamma_1 \Gamma_3 \langle \partial_1 G(x_1^n, x_3^n), \nu_1^n \rangle + \langle \partial_1 G(x_3^n, x_1^n), \nu_3^n \rangle
\end{aligned}$$

Lemma 3.7. $x_2^n \rightarrow p$ and $x_4^n \rightarrow p$ where $p \in \partial\Omega$ is from (3.4).

Proof. Suppose $|x_2^n - x_1^n| \geq c > 0$ and $|x_4^n - x_1^n| \geq c > 0$ along a subsequence, hence $\partial_1 G(x_i^n, x_j^n) = O(1)$ for $i \in \{1, 3\}, j \in \{2, 4\}$. Multiplying (3.6) by $\frac{2\pi d_1^n}{1 - \lambda_n}$, and using Lemma 3.2, (3.4) and $\lambda_n \rightarrow -1$, we obtain the contradiction:

$$\begin{aligned}
0 &= \Gamma_1^2 + \Gamma_3^2 \frac{d_1^n}{d_3^n} + 2 \frac{1 + \lambda_n}{1 - \lambda_n} \Gamma_1 \Gamma_3 d_1^n (d_1^n + d_3^n) \left(\frac{1}{|\bar{x}_1^n - x_3^n|^2} + \frac{1}{|\bar{x}_3^n - x_1^n|^2} \right) + o(1) \\
&\rightarrow \Gamma_1^2 + \Gamma_3^2.
\end{aligned}$$

Therefore we may assume that $x_2^n \rightarrow p$. Suppose $|x_1^n - x_4^n| \geq c > 0$ along a subsequence, hence $\partial_1 G(x_i^n, x_4^n) = O(1)$ for $i \in \{1, 3\}$. As above we multiply (3.6) by $\frac{2\pi d_1^n}{1 - \lambda_n}$ and obtain:

$$\begin{aligned}
0 &= \Gamma_1^2 + \Gamma_3^2 \frac{d_1^n}{d_3^n} + 2|\Gamma_1 \Gamma_2| d_1^n \left(\frac{\langle x_1^n - x_2^n, \nu_1^n \rangle}{|x_1^n - x_2^n|^2} - \frac{\langle x_1^n - \bar{x}_2^n, \nu_1^n \rangle}{|x_1^n - \bar{x}_2^n|^2} \right) \\
&\quad + 2|\Gamma_3 \Gamma_2| d_1^n \left(\frac{\langle x_3^n - x_2^n, \nu_3^n \rangle}{|x_3^n - x_2^n|^2} - \frac{\langle x_3^n - \bar{x}_2^n, \nu_3^n \rangle}{|x_3^n - \bar{x}_2^n|^2} \right) + o(1)
\end{aligned}$$

Passing to the limit now implies:

$$(3.7) \quad \Gamma_1^2 + \Gamma_3^2 + 2|\Gamma_2|(|\Gamma_1| + |\Gamma_3|)\alpha_1 \left(\alpha_1 - \alpha_2 - \frac{\alpha_1 + \alpha_2}{1 + 4\alpha_1\alpha_2} \right) = 0.$$

We used Lemma 3.6 for this computation. Observe that (3.7) implies $\alpha_1, \alpha_2 > 0$.

We also have

$$\begin{aligned}
0 &= \langle \partial_{x_2} (H(x^n) - \lambda_n \Phi(x^n)), \nu_2^n \rangle \\
&= (1 - \lambda_n) \left(\frac{\Gamma_2^2}{2\pi d_2^n} - 2|\Gamma_1 \Gamma_2| \langle \partial_1 G(x_2^n, x_1^n), \nu_2^n \rangle - 2|\Gamma_2 \Gamma_3| \langle \partial_1 G(x_2^n, x_3^n), \nu_2^n \rangle \right) \\
&\quad + 2(1 + \lambda_n) \Gamma_2 \Gamma_4 \langle \partial_1 G(x_2^n, x_4^n), \nu_2^n \rangle.
\end{aligned}$$

Since we know $x_1^n, x_2^n, x_3^n \rightarrow p$ and since we are assuming $|x_1^n - x_4^n| \geq c > 0$ we have $\partial_1 G(x_2^n, x_4^n) = O(1)$. Therefore multiplying the above equation by $\frac{2\pi d_2^n}{1-\lambda_n}$ we obtain as before

$$0 = \Gamma_2^2 + 2|\Gamma_1\Gamma_2|d_2^n \left(\frac{\langle x_2^n - x_1^n, \nu_2^n \rangle}{|x_2^n - x_1^n|^2} - \frac{\langle x_2^n - \bar{x}_1^n, \nu_2^n \rangle}{|x_2^n - \bar{x}_1^n|^2} \right) \\ + 2|\Gamma_2\Gamma_3|d_2^n \left(\frac{\langle x_2^n - x_3^n, \nu_2^n \rangle}{|x_2^n - x_3^n|^2} - \frac{\langle x_2^n - \bar{x}_3^n, \nu_2^n \rangle}{|x_2^n - \bar{x}_3^n|^2} \right) + o(1).$$

Again we pass to the limit and deduce:

$$(3.8) \quad \Gamma_2^2 + 2|\Gamma_2|(|\Gamma_1| + |\Gamma_3|)\alpha_2 \left(\alpha_2 - \alpha_1 - \frac{\alpha_1 + \alpha_2}{1 + 4\alpha_1\alpha_2} \right) = 0.$$

As before we used Lemma 3.6 for this computation. We need one more equation which comes from

$$0 = \langle \partial_{x_2} (H(x^n) - \lambda_n \Phi(x^n)), x_2^n - x_1^n \rangle \\ = (1 - \lambda_n) \left(\frac{\Gamma_2^2 \langle \nu_2^n, x_2^n - x_1^n \rangle}{2\pi d_2^n} - 2|\Gamma_1\Gamma_2| \langle \partial_1 G(x_2^n, x_1^n), x_2^n - x_1^n \rangle \right. \\ \left. - 2|\Gamma_2\Gamma_3| \langle \partial_1 G(x_2^n, x_3^n), x_2^n - x_1^n \rangle \right) + 2(1 + \lambda_n)\Gamma_2\Gamma_4 \langle \partial_1 G(x_2^n, x_4^n), x_2^n - x_1^n \rangle.$$

Since $\partial_1 G(x_2^n, x_4^n) = O(1)$ we get

$$0 = \Gamma_2^2 \frac{\langle \nu_2^n, x_2^n - x_1^n \rangle}{d_2^n} - 2|\Gamma_1\Gamma_2| \left(-\frac{\langle x_2^n - x_1^n, x_2^n - x_1^n \rangle}{|x_2^n - x_1^n|^2} + \frac{\langle x_2^n - \bar{x}_1^n, x_2^n - x_1^n \rangle}{|x_2^n - \bar{x}_1^n|^2} \right) \\ - 2|\Gamma_2\Gamma_3| \left(-\frac{\langle x_2^n - x_3^n, x_2^n - x_1^n \rangle}{|x_2^n - x_3^n|^2} + \frac{\langle x_2^n - \bar{x}_3^n, x_2^n - x_1^n \rangle}{|x_2^n - \bar{x}_3^n|^2} \right) + o(1).$$

Passing to the limit yields

$$(3.9) \quad \Gamma_2^2 \left(1 - \frac{\alpha_1}{\alpha_2} \right) + 4|\Gamma_2|(|\Gamma_1| + |\Gamma_3|)\alpha_1 \frac{\alpha_1 + \alpha_2}{1 + 4\alpha_1\alpha_2} = 0.$$

The system (3.7), (3.8), (3.9) has no solutions because $\alpha_2 \cdot (3.7) + \alpha_1 \cdot (3.8) + \alpha_2 \cdot (3.9)$ leads to $(\Gamma_1^2 + \Gamma_2^2 + \Gamma_3^2)\alpha_2 = 0$ which contradicts $\Gamma_i \neq 0, \alpha_2 > 0$. \square

Now we use (3.6) again. The same arguments as in the derivation of (3.7) lead to

$$(3.10) \quad \Gamma_1^2 + \Gamma_3^2 + 2|\Gamma_2|(|\Gamma_1| + |\Gamma_3|)\alpha_1 \left(\alpha_1 - \alpha_2 - \frac{\alpha_1 + \alpha_2}{1 + 4\alpha_1\alpha_2} \right) \\ + 2|\Gamma_4|(|\Gamma_1| + |\Gamma_3|)\beta_1 \left(\beta_1 - \beta_2 - \frac{\beta_1 + \beta_2}{1 + 4\beta_1\beta_2} \right) = 0.$$

The additional term involving β_1, β_2 comes from the fact that $x_4^n \rightarrow p$. In the derivation of (3.7) we assumed $|x_4^n - x_1^n| \geq c > 0$. This implies $\beta_1 = 0$, hence (3.7) is a special case of (3.10). We need to distinguish two cases:

CASE 1: $\liminf \frac{|x_2^n - x_4^n|}{d_2^n} = 0$

CASE 2: $\liminf \frac{|x_2^n - x_4^n|}{d_2^n} > 0$

In CASE 1, after passing to a subsequence we may assume that $|x_2^n - x_4^n| = o(d_2^n)$. This implies $\frac{d_2^n}{d_4^n} \rightarrow 1, \beta_1 = \alpha_1$ and $\beta_2 = \alpha_2$. Therefore (3.10) reduces to

$$(3.11) \quad \Gamma_1^2 + \Gamma_3^2 + 2(|\Gamma_2| + |\Gamma_4|)(|\Gamma_1| + |\Gamma_3|)\alpha_1 \left(\alpha_1 - \alpha_2 - \frac{\alpha_1 + \alpha_2}{1 + 4\alpha_1\alpha_2} \right) = 0.$$

Moreover, in CASE 1 (3.3) is also satisfied for $I = \{2, 4\}$. Thus we obtain

$$(3.12) \quad \Gamma_2^2 + \Gamma_4^2 + 2(|\Gamma_1| + |\Gamma_3|)(|\Gamma_2| + |\Gamma_4|)\alpha_2 \left(\alpha_2 - \alpha_1 - \frac{\alpha_1 + \alpha_2}{1 + 4\alpha_1\alpha_2} \right) = 0$$

in the same way as (3.11). We need one more equation which comes from

$$0 = \langle \partial_{x_2} (H(x^n) - \lambda_n \Phi(x^n)), \nu_2^n \rangle + \langle \partial_{x_4} (H(x^n) - \lambda_n \Phi(x^n)), \nu_4^n \rangle$$

Similar computations as before lead to

$$(3.13) \quad (\Gamma_2^2 + \Gamma_4^2) \left(1 - \frac{\alpha_1}{\alpha_2} \right) + 2(|\Gamma_2| + |\Gamma_4|)(|\Gamma_1| + |\Gamma_3|)2\alpha_1 \frac{\alpha_1 + \alpha_2}{1 + 4\alpha_1\alpha_2} = 0$$

Now $\alpha_2 \cdot (3.11) + \alpha_1 \cdot (3.12) + \alpha_2 \cdot (3.13)$ leads to $(\Gamma_1^2 + \Gamma_2^2 + \Gamma_3^2 + \Gamma_4^2)\alpha_2 = 0$ which contradicts $\Gamma_i \neq 0$, $\alpha_2 > 0$.

In CASE 2 we have $|x_2^n - x_4^n| \geq cd_2^n$ and $|x_2^n - x_4^n| \geq cd_4^n$. This implies

$$\langle \partial_1 G(x_2^n, x_4^n), x_2^n - x_1^n \rangle = O(1) = \langle \partial_1 G(x_4^n, x_2^n), x_2^n - x_1^n \rangle.$$

Then the equation

$$\langle \partial_{x_2} (H(x^n) - \lambda_n \Phi(x^n)), x_2^n - x_1^n \rangle = 0$$

leads to

$$(3.14) \quad \Gamma_2^2 \left(1 - \frac{\alpha_1}{\alpha_2} \right) + 2|\Gamma_2|(|\Gamma_1| + |\Gamma_3|)2\alpha_1 \frac{\alpha_1 + \alpha_2}{1 + 4\alpha_1\alpha_2} = 0.$$

Analogously, the equation

$$\langle \partial_{x_4} (H(x^n) - \lambda_n \Phi(x^n)), x_4^n - x_1^n \rangle = 0$$

leads to

$$(3.15) \quad \Gamma_4^2 \left(1 - \frac{\beta_1}{\beta_2} \right) + 2|\Gamma_4|(|\Gamma_1| + |\Gamma_3|)2\beta_1 \frac{\beta_1 + \beta_2}{1 + 4\beta_1\beta_2} = 0.$$

Finally the equations

$$\langle \partial_{x_4} (H(x^n) - \lambda_n \Phi(x^n)), \nu_4^n \rangle = 0$$

and

$$\langle \partial_{x_2} (H(x^n) - \lambda_n \Phi(x^n)), \nu_4^n \rangle = 0$$

lead, respectively, to

$$(3.16) \quad \Gamma_2^2 + 2|\Gamma_2|(|\Gamma_1| + |\Gamma_3|)\alpha_2 \left(\alpha_2 - \alpha_1 - \frac{\alpha_1 + \alpha_2}{1 + 4\alpha_1\alpha_2} \right) = 0.$$

and

$$(3.17) \quad \Gamma_4^2 + 2|\Gamma_4|(|\Gamma_1| + |\Gamma_3|)\alpha_2 \left(\beta_2 - \beta_1 - \frac{\beta_1 + \beta_2}{1 + 4\beta_1\beta_2} \right) = 0.$$

Now the sum $\alpha_2\beta_2 \cdot (3.10) + \alpha_2\beta_2 \cdot (3.14) + \alpha_1\beta_2 \cdot (3.15) + \alpha_2\beta_1 \cdot (3.16) + \alpha_2\beta_2 \cdot (3.17)$ leads to $(\Gamma_1^2 + \Gamma_2^2 + \Gamma_3^2 + \Gamma_4^2)\alpha_2\beta_2 = 0$ which as before contradicts $\Gamma_i \neq 0$, $\alpha_2, \beta_2 > 0$. This concludes the proof of Proposition 3.1.

4 Proof of Theorems 2.1-2.3

Proof of Theorem 2.1. There exists a compact subset $K_0 \subset \mathcal{F}_2\Omega$ such that $\text{cat}(K_0) = \text{cat}(\mathcal{F}_2\Omega)$. Observe that

$$H(x) = \Gamma_1^2 h(x_1) + \Gamma_2^2 h(x_2) + \Gamma_1 \Gamma_2 G(x_1, x_2) + f(x) \rightarrow -\infty \quad \text{as } x \rightarrow \partial \mathcal{F}_2\Omega$$

because $\Gamma_1 \Gamma_2 < 0$, $f(x) = O(1)$, and assumption (A2). Therefore $H^{\geq a} = \{x \in \mathcal{F}_2\Omega : H(x) \geq a\}$ is compact for any $a \in \mathbb{R}$. Now we choose $a < \min H(K_0)$, set $\delta := \frac{1}{2}(\min H(K_0) - a)$, and consider F on the compact manifold $K = H^{\geq a}$ with boundary $B = H^{-1}(a)$. Since $\min F(K) > \max F(B)$ standard critical point theory yields that a function $F \in \mathcal{C}^1(\mathcal{F}_N\Omega)$ with $\|F|_K - H|_K\| < \delta$ has at least $\text{cat}(\mathcal{F}_2\Omega)$ critical points in K . This proves a).

Part b) follows similarly upon passing to the quotient $\mathcal{F}_2(\Omega)/\langle (x_1, x_2) \sim (x_2, x_1) \rangle$. Finally, c) is obvious. \square

The proof of Theorem 2.2 and of Theorem 2.3 will be based on a linking argument. In the sequel N will be either 3 or 4. Suppose there exists a (sequentially) compact topological space S , a continuous map $\gamma_0 : S \rightarrow \mathcal{F}_N\Omega$, and a subset $\mathcal{L} \subset \mathcal{F}_N\Omega$ such that

$$(4.1) \quad \sup_{x \in \mathcal{L}} H(x) < \infty,$$

and

$$(4.2) \quad \gamma \text{ is homotopic to } \gamma_0 \implies \gamma(S) \cap \mathcal{L} \neq \emptyset.$$

As usual, γ being homotopic to γ_0 means that there exists a continuous deformation $H : S \times [0, 1] \rightarrow \mathcal{F}_N\Omega$ with $H(\zeta, 0) = \gamma_0(\zeta)$ and $H(\zeta, 1) = \gamma(\zeta)$ for all $\zeta \in S$. We shall prove that if a function $F \in \mathcal{C}^1(\mathcal{F}_N\Omega, \mathbb{R})$ is close to H on compact sets then it has a critical point. In order to express the closeness we choose $a < \min_{\zeta \in S} H(\gamma_0(\zeta))$ and $b > \sup_{x \in \mathcal{L}} H(x)$. Let M_0 be as in Proposition 3.1 for these values $a < b$. By Sard's theorem we may assume that $-M_0$ is a regular value of Φ . Since S is sequentially compact we may also assume that $-M_0 < \inf_{\zeta \in S} \Phi(\zeta)$. Setting

$$V_\Omega(x) := \nabla H(x) - \frac{\langle \nabla H(x), \nabla \Phi(x) \rangle}{|\nabla \Phi(x)|^2} \nabla \Phi(x)$$

Proposition 3.1 implies

$$a \leq H(x) \leq b, \Phi(x) = -M_0, \langle \nabla H(x), \nabla \Phi(x) \rangle \leq 0 \implies V_\Omega(x) \neq 0.$$

Observe that $D := \{x \in \mathcal{F}_N\Omega : \Phi(x) \geq -M_0\}$ is a compact manifold with smooth boundary $\partial D = \Phi^{-1}(-M_0)$. We also define

$$D_a^b := \{x \in D : a \leq H(x) \leq b\} = \{x \in \mathcal{F}_N\Omega : \Phi(x) \geq -M_0, a \leq H(x) \leq b\}.$$

Now we choose $\varepsilon > 0$ satisfying

$$a + 2\varepsilon < \min_{\zeta \in S} H(\gamma_0(\zeta)) \leq \sup_{x \in \mathcal{L}} H(x) < b - 2\varepsilon$$

and

$$(4.3) \quad \varepsilon < \frac{1}{2} \min\{|V_\Omega(x)| : a \leq H(x) \leq b, \Phi(x) = -M_0, \langle \nabla H(x), \nabla \Phi(x) \rangle \leq 0\}.$$

Proposition 4.1. Suppose $F \in \mathcal{C}^1(\mathcal{F}_N\Omega, \mathbb{R})$ satisfies

$$(4.4) \quad |F(x) - H(x)| \leq \varepsilon \quad \text{if } x \in K := D_a^b,$$

and

$$(4.5) \quad |\nabla F(x) - \nabla H(x)| \leq \varepsilon \quad \text{if } x \in D_a^b \cap \partial D.$$

Then F has a critical point in $K = D_a^b$.

Clearly (4.4) requires F to be \mathcal{C}^0 -close to H on the compact set D_a^b , and (4.5) requires F to be \mathcal{C}^1 -close to H on the compact set $D_a^b \cap \partial D$.

Proof. We assume that F has no critical value in D_a^b . First we define a continuous map $V_0 : D_a^b \cap \partial D \rightarrow \mathbb{R}^{2N}$ by setting:

$$V_0(x) := \begin{cases} \nabla F(x) - \frac{\langle \nabla F(x), \nabla \Phi(x) \rangle}{|\nabla \Phi(x)|^2} \nabla \Phi(x) & \text{if } \langle \nabla F(x), \nabla \Phi(x) \rangle \leq 0; \\ \nabla F(x) & \text{else.} \end{cases}$$

Clearly we have

$$(4.6) \quad \langle V_0(x), \nabla \Phi(x) \rangle \geq 0 \quad \text{for all } x \in D_a^b \cap \partial D,$$

hence $V_0(x)$ is either tangent to ∂D at x or points inside D . Using (4.3) and (4.5) it is easy to check that

$$(4.7) \quad \langle \nabla F(x), V_0(x) \rangle > 0 \quad \text{if } x \in D_a^b \cap \partial D.$$

Next we extend this vector field to all of $\mathcal{F}_N\Omega$. In order to do this we first choose a relatively open tubular neighborhood $\partial D \subset \mathcal{O} \subset D$ of ∂D and a diffeomorphism $\chi = (\chi_1, \chi_2) : \mathcal{O} \rightarrow \partial D \times [0, 1]$ such that $\chi(x) = (x, 0)$ for $x \in \partial D$. Then we define for $0 < \delta < 1$ a map $V_1 : D_a^b \rightarrow \mathbb{R}^{2N}$ by setting

$$V_1(x) := \begin{cases} \frac{\delta - \chi_2(x)}{\delta} V_0(\chi_1(x)) + \frac{\chi_2(x)}{\delta} \nabla F(x) & \text{if } x \in D_a^b \cap \mathcal{O}, \chi_2(x) \leq \delta; \\ \nabla F(x) & \text{if } x \in D_a^b \cap \mathcal{O}, \chi_2(x) > \delta, \text{ or } x \in D_a^b \setminus \mathcal{O}. \end{cases}$$

Observe that V_1 is continuous and coincides with V_0 on $D_a^b \cap \partial D$. Therefore, if $\delta > 0$ is small (4.7) implies that

$$(4.8) \quad \langle \nabla F(x), V_1(x) \rangle > 0 \quad \text{if } x \in D_a^b.$$

Here we also used that F has no critical point in D_a^b . We fix such a $\delta > 0$. Then we replace the continuous vector field $V_1 : D_a^b \rightarrow \mathbb{R}^{2N}$ by a Lipschitz continuous vector field $V_F : D_a^b \rightarrow \mathbb{R}^{2N}$ such that (4.6) and (4.8) continue to hold for V_F instead of V_0, V_1 . Finally we extend the vector field $V_F : D_a^b \rightarrow \mathbb{R}^{2N}$ to a Lipschitz continuous vector field $V_F : \mathcal{F}_N\Omega \rightarrow \mathbb{R}^{2N}$ such that $V_F(x) = 0$ outside a neighborhood of D_a^b , and such that $\langle \nabla F(x), V_F(x) \rangle \geq 0$ for all $x \in \mathcal{F}_N\Omega$. As a consequence, V_F defines a global flow $\varphi : \mathcal{F}_N\Omega \times \mathbb{R} \rightarrow \mathcal{F}_N\Omega$ which satisfies:

$$(4.9) \quad x \in D, a \leq H(\varphi(x, t)) \leq b \text{ for } 0 \leq t \leq T \quad \implies \quad \varphi(x, T) \in D$$

and

$$(4.10) \quad \begin{cases} x \in D, a \leq H(\varphi(x, t)) \leq b \text{ for all } t \geq 0, \\ \implies \varphi(x, t_n) \rightarrow \bar{x} \text{ for some sequence } t_n \rightarrow \infty, \nabla F(\bar{x}) = 0. \end{cases}$$

Now we argue as follows. By (4.2) for each $n \in \mathbb{N}$ there exists $\zeta_n \in S$ such that $\varphi(\gamma_0(\zeta_n), n) \in \mathcal{L}$, hence $a \leq H(\gamma_0(\zeta_n), n) \leq b$. Since S is sequentially compact we have $\zeta_n \rightarrow \zeta \in S$ along a subsequence. It follows that $x := \gamma_0(\zeta) \in D_a^b$ satisfies $\varphi(x, t) \in D_a^b$ for all $t \geq 0$. Now the existence of a critical point of F in D_a^b follows from (4.10). \square

In the proof of Theorems 2.2 and 2.3 the set \mathcal{L} will be

$$\mathcal{L}_3(\Omega) := \{x \in \mathcal{F}_3\Omega : x_1 - x_2 + r(x_3 - x_2) = 0 \text{ for some } r > 0\},$$

in case $N = 3$, and the set

$$\mathcal{L}_4(\Omega) := \{x \in \mathcal{F}_4\Omega : x_1 - x_2 + r(x_3 - x_2) = 0, x_2 - x_3 + s(x_4 - x_3) = 0 \text{ for some } r, s > 0\}.$$

in case $N = 4$, as in [6]. So we need to bound H on these sets.

Lemma 4.2. *a) Suppose $N = 3$ and (2.3) holds. Then $\sup_{\mathcal{L}_3\Omega} H < \infty$.*

b) Suppose $N = 4$ and (2.3) and (2.4) hold. Then $\sup_{\mathcal{L}_4\Omega} H < \infty$.

Proof. We shall prove that if $x^n \in \mathcal{L}_N\Omega$ is such that $x^n \rightarrow \partial\mathcal{L}_N\Omega$, then $H(x^n) \rightarrow -\infty$. Set $d_i^n := \text{dist}(x_i^n, \partial\Omega)$. As in Section 3 we drop the notation $n \rightarrow \infty$ from all limits, in particular for the terms $O(1)$ and $o(1)$. Without loss of generality we may assume that $\Gamma_i = (-1)^i k_i$ with $k_i > 0$.

a) The Hamiltonian has the form

$$H(x) = \sum_{i=1}^3 k_i^2 h(x_i) - 2k_1 k_2 G(x_1, x_2) + 2k_1 k_3 G(x_1, x_3) - 2k_2 k_3 G(x_2, x_3) + f(x),$$

and assumption (2.3) reads as

$$(4.11) \quad k_1 k_2 + k_2 k_3 - k_1 k_3 > 0.$$

Observe that if $x \in \mathcal{L}_3\Omega$ then

$$(4.12) \quad |x_1 - x_3| > \max\{|x_1 - x_2|, |x_2 - x_3|\}.$$

If $d_i^n \geq c > 0$ for every i then $h(x_i^n), g(x_i^n, x_j^n) = O(1)$, and $|x_i^n - x_j^n| \rightarrow 0$ for at least one $i \neq j$. Then (4.11) and (4.12) imply

$$\begin{aligned} H(x^n) &= \sum_{i=1}^3 k_i^2 h(x_i^n) - 2k_1 k_2 g(x_1^n, x_2^n) + 2k_1 k_3 g(x_1^n, x_3^n) - 2k_2 k_3 g(x_2^n, x_3^n) \\ &\quad + \frac{1}{\pi} \log \frac{|x_1^n - x_2^n|^{k_1 k_2} |x_2^n - x_3^n|^{k_2 k_3}}{|x_1^n - x_3^n|^{k_1 k_3}} + f(x^n) \\ &= \frac{1}{\pi} \log \frac{|x_1^n - x_2^n|^{k_1 k_2} |x_2^n - x_3^n|^{k_2 k_3}}{|x_1^n - x_3^n|^{k_1 k_3}} + O(1) \rightarrow -\infty \end{aligned}$$

Thus we may assume from now on that $d_i^n \rightarrow 0$ for some i . If in addition $|x_j^n - x_\ell^n| \geq c > 0$ for every $j \neq \ell$ then $H(x^n) \rightarrow -\infty$ by (A2) and because $f(x^n) = O(1)$. It follows that we only need to consider the case where $|x_j^n - x_\ell^n| \rightarrow 0$ for some j and ℓ . Observe that if only one of $|x_1^n - x_2^n| \rightarrow 0$ or $|x_2^n - x_3^n| \rightarrow 0$ hold then (A1) and (A2) immediately imply $H(x^n) \rightarrow -\infty$. Therefore we may assume that $|x_1^n - x_3^n| \rightarrow 0$ and $d_i^n \rightarrow 0$ for some i , hence $d_i^n \rightarrow 0$ for every i because of (4.12). Thus we are left with the following case:

$$|x_1^n - x_3^n| \rightarrow 0, d_i^n \rightarrow 0 \text{ for all } i.$$

If $\frac{|x_1^n - x_3^n|}{d_1^n} \geq c > 0$ Lemma 3.2 (iv) implies $G(x_1^n, x_3^n) = O(1)$, and the claim follows. Therefore it remains to consider the case $|x_1^n - x_3^n| = o(d_1^n)$, hence also $|x_1^n - x_3^n| = o(d_3^n)$. By (4.12) we also have

$|x_1^n - x_2^n| = o(d_1^n)$, and $d_i^n/d_j^n \rightarrow 1$ for all i, j . Furthermore we can deduce that $|x_1^n - \bar{x}_2^n|, |x_3^n - \bar{x}_2^n| \geq d_2^n = d_1^n(1 + o(1))$, and $|x_1^n - \bar{x}_3^n| \leq cd_1^n$. Now (i) and (iii) of Lemma 3.2 yield

$$\begin{aligned} 2\pi H(x^n) &= \log \left((d_1^n)^{k_1^2} (d_2^n)^{k_2^2} (d_3^n)^{k_3^2} \frac{|x_1^n - x_2^n|^{2k_1 k_2} |x_1^n - \bar{x}_3^n|^{2k_1 k_3} |x_2^n - x_3^n|^{2k_2 k_3}}{|x_1^n - \bar{x}_2^n|^{2k_1 k_2} |x_1^n - x_3^n|^{2k_1 k_3} |\bar{x}_2^n - x_3^n|^{2k_2 k_3}} \right) + O(1) \\ &\leq \log \left((d_1^n)^{k_1^2} (d_2^n)^{k_2^2} (d_3^n)^{k_3^2} \frac{|x_1^n - x_3^n|^{2k_1 k_2 + 2k_2 k_3 - 2k_1 k_3} |x_1^n - \bar{x}_3^n|^{2k_1 k_3}}{(d_2^n)^{2k_1 k_2 + 2k_2 k_3}} \right) + O(1) \\ &\leq \log \left(c(d_1^n)^{k_1^2} (d_2^n)^{k_2^2} (d_3^n)^{k_3^2} \left(\frac{|x_1^n - x_3^n|}{d_1^n} \right)^{2k_1 k_2 + 2k_2 k_3 - 2k_1 k_3} \right) + O(1) \rightarrow -\infty, \end{aligned}$$

for some constant $c > 0$. For the convergence we used assumption (4.11) and $|x_1^n - x_3^n| = o(d_1^n)$.

b) Here the Hamiltonian has the form

$$\begin{aligned} H(x) &= \sum_{i=1}^4 k_i^2 h(x_i) - 2k_1 k_2 G(x_1, x_2) + 2k_1 k_3 G(x_1, x_3) - 2k_1 k_4 G(x_1, x_4) \\ &\quad - 2k_2 k_3 G(x_2, x_3) + 2k_2 k_4 G(x_2, x_4) - 2k_3 k_4 G(x_3, x_4) + f(x). \end{aligned}$$

Assumption (2.3) implies

$$(4.13) \quad k_1 k_2 + k_2 k_3 - k_1 k_3 > 0, \quad k_2 k_3 + k_3 k_4 - k_2 k_4 > 0,$$

and assumption (2.4) implies

$$(4.14) \quad k_1(k_2 + k_4 - k_3) > 0, \quad k_4(k_1 + k_3 - k_2) > 0.$$

For $x \in \mathcal{L}_4\Omega$ there holds

$$(4.15) \quad \begin{aligned} |x_1 - x_3| &> \max\{|x_1 - x_2|, |x_2 - x_3|\}, \quad |x_2 - x_4| > \max\{|x_2 - x_3|, |x_3 - x_4|\} \\ |x_1 - x_4| &> \max\{|x_1 - x_3|, |x_2 - x_4|\}. \end{aligned}$$

If $|x_j^n - x_\ell^n| \geq c > 0$ for every $j \neq \ell$ then $d_i^n \rightarrow 0$ for some i and $H(x^n) \rightarrow -\infty$ as a consequence of (A1) and (A2). If $|x_j^n - x_\ell^n| \rightarrow 0$ for some $j \neq \ell$ then the only case we have to check is when $|x_1^n - x_4^n| \rightarrow 0$ because all the other cases can be treated as in the proof of a). In this case, if $d_i^n \geq c > 0$ for every i we have

$$H(x^n) = \frac{1}{\pi} \log \frac{|x_1^n - x_2^n|^{k_1 k_2} |x_1^n - x_4^n|^{k_1 k_4} |x_2^n - x_3^n|^{k_2 k_3} |x_3^n - x_4^n|^{k_3 k_4}}{|x_1^n - x_3^n|^{k_1 k_3} |x_2^n - x_4^n|^{k_2 k_4}} + O(1) \rightarrow -\infty,$$

because for some $c > 0$

$$\begin{aligned} &\frac{|x_1^n - x_2^n|^{k_1 k_2} |x_1^n - x_4^n|^{k_1 k_4} |x_2^n - x_3^n|^{k_2 k_3} |x_3^n - x_4^n|^{k_3 k_4}}{|x_1^n - x_3^n|^{k_1 k_3} |x_2^n - x_4^n|^{k_2 k_4}} \\ &\leq c \left(|x_1^n - x_3^n|^{k_1 k_4} + |x_3^n - x_4^n|^{k_1 k_4} \right) \frac{|x_1^n - x_2^n|^{k_1 k_2} |x_2^n - x_3^n|^{k_2 k_3} |x_3^n - x_4^n|^{k_3 k_4}}{|x_1^n - x_3^n|^{k_1 k_3} |x_2^n - x_4^n|^{k_2 k_4}} \\ &\leq c |x_1^n - x_3^n|^{k_1 k_2 + k_1 k_4 - k_1 k_3} |x_2^n - x_4^n|^{k_2 k_3 + k_3 k_4 - k_2 k_4} \\ &\quad + c |x_1^n - x_3^n|^{k_1 k_2 + k_2 k_3 - k_1 k_3} |x_2^n - x_4^n|^{k_1 k_4 + k_3 k_4 - k_2 k_4} \\ &\rightarrow 0. \end{aligned}$$

Here we used (4.13), (4.14), and (4.15).

It remains to consider the case when $|x_1^n - x_4^n| \rightarrow 0$ and $d_i^n \rightarrow 0$ for some i which implies $d_i^n \rightarrow 0$ for every i by (4.15). If

$$(4.16) \quad \frac{|x_1^n - x_3^n|}{d_1^n} \geq c > 0$$

and

$$(4.17) \quad \frac{|x_2^n - x_4^n|}{d_2^n} \geq c > 0$$

Lemma 3.2 (iv) implies $G(x_1^n, x_3^n) = O(1)$ and $G(x_2^n, x_4^n) = O(1)$ and the claim follows. If only one of (4.16), (4.17) is true we argue as in a).

Finally, we are left with the case $|x_1^n - x_3^n| = o(d_1^n)$ and $|x_2^n - x_4^n| = o(d_2^n)$. In this case, it is easy to check that

$$(4.18) \quad |x_1^n - x_4^n| = o(d_1^n) \quad \text{and} \quad \frac{d_i^n}{d_j^n} \rightarrow 1.$$

Setting $q_{i,j}^n := \frac{|x_i^n - x_j^n|}{|x_i^n - \bar{x}_j^n|}$, Lemma 3.2 (i), (ii) yields

$$2\pi H(x^n) = \log(d_1^n)^{k_1^2} (d_2^n)^{k_2^2} (d_3^n)^{k_3^2} (d_4^n)^{k_4^2} + \log \frac{q_{1,2}^n q_{1,4}^n q_{2,3}^n q_{3,4}^n}{q_{1,3}^n q_{2,4}^n} + O(1).$$

From (4.14), (4.15), (4.18), we deduce $d_j^n \leq |x_i^n - \bar{x}_j^n| \leq 3d_j^n$, hence $q_{1,2}^n \leq \frac{c|x_1^n - x_3^n|}{d_1^n}$ for some $c > 0$, and similarly for the other $q_{i,j}^n$. Using this and $|x_1^n - x_4^n| \leq |x_1^n - x_3^n| + |x_2^n - x_4^n|$ we obtain

$$\begin{aligned} \frac{q_{1,2}^n q_{1,4}^n q_{2,3}^n q_{3,4}^n}{q_{1,3}^n q_{2,4}^n} &\leq c \left(\frac{|x_1^n - x_3^n|}{d_1^n} \right)^{2k_1 k_2 - 2k_1 k_3 + 2k_1 k_4} \left(\frac{|x_2^n - x_4^n|}{d_2^n} \right)^{2k_2 k_3 + 2k_3 k_4 - 2k_2 k_4} \\ &\quad + c \left(\frac{|x_1^n - x_3^n|}{d_1^n} \right)^{2k_1 k_2 + 2k_2 k_3 - 2k_1 k_2} \left(\frac{|x_2^n - x_4^n|}{d_2^n} \right)^{2k_1 k_4 + 2k_3 k_4 - 2k_2 k_4} \\ &\rightarrow 0 \end{aligned}$$

Thus also in this case $H(x^n) \rightarrow -\infty$. \square

Proof of Theorem 2.2. We recall the linking from [6]. We assume without loss of generality that $0 \in \Omega$ and fix $\rho > 0$ such that the closed ball $B(0, 2\rho) \subset \Omega$. Using complex notation for the elements of $\Omega \subset \mathbb{R}^2 = \mathbb{C}$, we set

$$(4.19) \quad \gamma_0 : S^1 = \{\zeta \in \mathbb{C} : |\zeta| = 1\} \rightarrow \mathcal{F}_3\Omega, \quad \gamma_0(\zeta) := (\rho\zeta, 0, 2\rho).$$

Then (4.2) holds for $S = S^1$, γ_0 from (4.19), and $\mathcal{L} = \mathcal{L}_3(\Omega)$. This has been proved in [6, Lemma 6.2]. It follows that a \mathcal{C}^1 -function $F : \mathcal{F}_3\Omega \rightarrow \mathbb{R}$ which is \mathcal{C}^1 -close to H in the sense of Proposition 4.1 has a critical point. This proves part a) of Theorem 2.2, part b) is proved easily. \square

Proof of Theorem 2.3. For $N = 4$ vortices we set

$$(4.20) \quad \gamma_0 : S^1 \times S^1 \rightarrow \mathcal{F}_4\Omega, \quad \gamma_0(\zeta_1, \zeta_2) := (\rho\zeta_1, 0, 3\rho, 3\rho + \rho\zeta_2).$$

It has been proved in [6, Lemma 7.2] that (4.2) holds for $S = S^1 \times S^1$, γ_0 from (4.20), and $\mathcal{L} = \mathcal{L}_4(\Omega)$. As above it follows that a \mathcal{C}^1 -function $F : \mathcal{F}_4\Omega \rightarrow \mathbb{R}$ which is \mathcal{C}^1 -close to H in the sense of Proposition 4.1 has a critical point. \square

5 Proof of Theorem 2.4

Following [7] we prove Theorem 2.4 by constructing streamfunctions ψ_ε as solutions of the elliptic problem

$$(5.1) \quad \begin{cases} -\varepsilon^2 \Delta \psi = \sum_{i=1}^N f_i \left(\psi + \frac{\Gamma_i}{2\pi} \ln \varepsilon \right) & \text{in } \Omega; \\ \psi = \psi_0 & \text{on } \partial\Omega. \end{cases}$$

with $f_i(t) = t_+^p$ if $\Gamma_i > 0$, and $f_i(t) = -t_-^p$ if $\Gamma_i < 0$; here $t_\pm = \max\{\pm t, 0\}$ and $1 < p < \frac{N+2}{N-2}$.

Setting $u = \frac{2\pi}{|\ln \varepsilon|}(\psi - \psi_0)$ and $\delta = \varepsilon \left(\frac{2\pi}{|\ln \varepsilon|} \right)^{(p-1)/2}$ these are obtained as critical points of the functional $I : H_0^1(\Omega) \rightarrow \mathbb{R}$ defined by

$$(5.2) \quad I(u) = \frac{\delta^2}{2} \int_{\Omega} |\nabla u|^2 - \sum_{i=1}^N \int_{\Omega} F_i \left(u - \Gamma_i - \frac{2\pi\psi_0(x)}{|\ln \varepsilon|} \right)$$

with $F_i(t) = \int_0^t f_i(s) ds$. Choose $R > 0$ such that $\Omega \subset \subset B_R(0)$. For $a > 0$ let $W_{\delta,a}$ be the unique positive solution of

$$\begin{cases} -\delta^2 \Delta w = (w - a)_+^p & \text{in } B_R(0); \\ w = 0 & \text{on } \partial B_R(0), \end{cases}$$

and define $W_{\delta,x,a}(y) := W_{\delta,a}(y-x)$ for $x, y \in \Omega$. Finally, let $P : H_0^1(B_R(0)) \rightarrow H_0^1(\Omega)$ be the orthogonal projection, hence $w = PW_{\delta,x,a}$ solves

$$\begin{cases} -\delta^2 \Delta w = (W_{\delta,x,a} - a)_+^p & \text{in } \Omega; \\ w = 0 & \text{on } \partial\Omega. \end{cases}$$

Now in order to obtain a solution of (5.1), for $x \in \mathcal{F}_N \Omega$ and $a_i > 0$ one makes the ansatz

$$u = \sum_{i=1}^N (\text{sign } \Gamma_i) PW_{\delta,x_i,a_i} + w_\delta$$

with w_δ a small perturbation. Then a Lyapunov-Schmidt procedure yields $w_{\delta,x} \in H_0^1(\Omega)$ with $\|w_{\delta,x}\|_\infty = O(|\delta| \ln |\delta|^{(p-1)/2})$ and $a_{i,\delta}(x) > 0$ such that the following holds: If $x \in \mathcal{F}_N \Omega$ is a critical point of

$$F_\delta(x) := I \left(\sum_{i=1}^N (\text{sign } \Gamma_i) PW_{\delta,x_i,a_{i,\delta}(x)} + w_{\delta,x} \right)$$

then

$$(5.3) \quad u_\delta = \sum_{i=1}^N (\text{sign } \Gamma_i) PW_{\delta,x_i,a_{i,\delta}(x)} + w_{\delta,x}$$

is a critical point of I ; see [7, Section 3].

By [7, (4.2), (4.3)] there holds

$$F_\delta(x) = \alpha(\delta) + \beta(\delta) H_{KR}(x) + \chi_\delta(x)$$

where $\alpha(\delta)$ and $\beta(\delta)$ are independent of x , and χ_δ converges to 0 as $\delta \rightarrow 0$ uniformly in the \mathcal{C}^1 -norm on compact sets of $\mathcal{F}_N \Omega$.

Now Theorems 2.1–2.3 yield for $\delta > 0$ small critical points $x_\delta \in \mathcal{F}_N\Omega$ of F_δ such that $x_\delta \rightarrow x^*$ along a subsequence, where $x^* \in \mathcal{F}_N\Omega$ is a critical point of H_{KR} . As a consequence we obtain corresponding critical points u_δ of I as in (5.3), hence solutions v_δ of the Euler equation (1.3). That the scalar vorticity $\omega_\delta = \nabla \times v_\delta = -\Delta u_\delta$ concentrates near x^* follows as in [7] from the fact that $\Delta PW_{\delta, x_i, a_i, \delta}(x) = 0$ if $W_{\delta, x_i, a_i, \delta}(x) < a_i$.

References

- [1] H. Aref: *Point vortex dynamics: A classical mathematical playground*. J. Math. Phys. **48** (2007), 1–22
- [2] H. Aref, P. K. Newton, M. A. Strember, T. Tokieda, D. Vainchtein: *Vortex crystals*. Adv. Appl. Mech. **39** (2002), 1–79
- [3] C. Bandle, M. Flucher: *Harmonic radius and concentration of energy; hyperbolci radius and Liouville’s equations $\Delta U = e^U$ and $\Delta U = U^{\frac{n+2}{n-2}}$* . SIAM Reviews **38** (1996), 191–238.
- [4] D. Bartolucci, A. Pistoia: *Existence and qualitative properties of concentrating solutions for the sinh-Poisson equation*. IMA J. Appl. Math. **72** (2007), 706–729.
- [5] T. Bartsch, Q. Dai: *Periodic solutions of the N -vortex problem in planar domains*. Preprint.
- [6] T. Bartsch, A. Pistoia, T. Weth: *N -vortex equilibria for ideal fluids in bounded planar domains and new nodal solutions of the sinh-Poisson and the Lane-Emden-Fowler equations*. Comm. Math. Phys. **297** (2010), 653–687.
- [7] D. Cao, Z. Liu, J. Wei: *Regularization of Point Vortices Pairs for the Euler Equation in Dimension Two*. Arch. Rat. Mech. Anal. **212** (2014), 179–217.
- [8] D. Cao, Z. Liu, J. Wei: *Regularization of Point Vortices Pairs for the Euler Equation in Dimension Two, Part II*. arXiv:1208.5540.
- [9] M. del Pino, M. Kowalczyk, M. Musso: *Singular limits in Liouville-type equations*. Calc. Var. Part. Diff. Equ. **24** (2005), 47–81.
- [10] P. Esposito, M. Musso, A. Pistoia: *Concentrating solutions for a planar elliptic problem involving nonlinearities with large exponent*. J. Diff. Equ. **227** (2006), no. 1, 29–68.
- [11] P. Esposito, M. Musso, A. Pistoia: *On the existence and profile of nodal solutions for a two-dimensional elliptic problem with large exponent in nonlinearity*. J. London Math. Soc. (3) **94** (2007), 497–519.
- [12] P. Esposito, M. Grossi, A. Pistoia: *On the existence of blowing-up solutions for a mean field equation*. Ann. Inst. H. Poincaré, Anal. non lin. **22** (2005), 227–257,
- [13] M. Flucher, B. Gustafsson: *Vortex motion in two-dimensional hydromechanics*. Preprint in TRITA-MAT-1997-MA-02.
- [14] H. Helmholtz: *Über Integrale der hydrodynamischen Gleichungen, welche den Wirbelbewegungen entsprechen*. J. reine angew. Math. **55** (1858), 25–55
- [15] G. Kirchhoff: *Vorlesungen über mathematische Physik*. Teubner, Leipzig 1876
- [16] C. Kuhl: *Equilibria for the N -vortex problem in a general bounded domain*. Preprint 2014.
- [17] C. Kuhl: *Symmetric equilibria for the N -vortex problem*. Preprint 2014.

- [18] C. C. Lin: *On the motion of vortices in 2D I. Existence of the Kirchhoff-Routh function*. Proc. Nat. Acad. Sc. **27** (1941), 570–575.
- [19] C. C. Lin: *On the motion of vortices in 2D II. Some further properties on the Kirchhoff-Routh function*. Proc. Nat. Acad. Sc. **27** (1941), 575–577.
- [20] A. J. Majda, A. L. Bertozzi: *Vorticity and Incompressible Flow*. Cambridge University Press 2001.
- [21] C. Marchioro, M. Pulvirenti: *Mathematical Theory of Incompressible Nonviscous Fluids*. Applied Mathematical Sciences **96**, Springer-Verlag, New York 1994.
- [22] P. K. Newton: *The N-vortex problem*. Springer-Verlag, Berlin 2001.
- [23] E. J. Routh: *Some applications of conjugate functions*. Proc. London Mat. Soc. **12** (1881), 73–89.
- [24] P. G. Saffman: *Vortex dynamics*. Cambridge University Press, Cambridge 1992.

ADDRESS OF THE AUTHORS:

Thomas Bartsch	Angela Pistoia
Mathematisches Institut	Dipartimento di Metodi e Modelli Matematici
University of Giessen	Università di Roma "La Sapienza"
Arndtstr. 2	via A. Scarpa 16
35392 Giessen	00161 Roma
Germany	Italy
Thomas.Bartsch@math.uni-giessen.de	pistoia@dmmm.uniroma1.it