


# Synthesis Challenge AG Wegner

JLU Giessen

24.10.2013




↓  
1,2,3




A

- 1) LiHMDS (1.25 equiv),  $t\text{Bu}-\text{Me}_2\text{SiCl}$  (1.25 equiv),  $-78^\circ\text{C}$
- 2) PhNTf<sub>2</sub> (1.5 equiv), CsF (2.5 equiv), ( $\text{MeOCH}_2$ )<sub>2</sub>,
- 3) I (1.5 equiv),  $[\text{Pd}(\text{dppf})\text{Cl}_2]\cdot\text{CH}_2\text{Cl}_2$  (10 mol%),  $\text{Ph}_3\text{As}$  (10 mol%),  $\text{Cs}_2\text{CO}_3$



I




↓  
4,5

B


- 4) 9-BBN (1.1 equiv), THF, 0.8C to RT; then II (1.0 equiv),  $[\text{Pd}(\text{dppf})\text{Cl}_2]\cdot\text{CH}_2\text{Cl}_2$  (2.7 mol %), NaOH
- 5) PPTS (10 mol %), MeOH

please provide a detailed mechanism for step 4)



6

6)  $[\{Ir(cod)Cl\}_2]$  (3.2 mol%)  
 $Zn(OTf)_2$  (16 mol%), (*R*)-III  
(12.8 mol%)




7-10

7) DDQ (1.1 equiv), pH 7 buffer,  
8) DMP (1.5 equiv),  
9)  $NaClO_2$  (4.0 equiv),  $NaH_2PO_4$   
(6.0 equiv), 2-methyl-2-butene  
(70 equiv), tBuOH/H<sub>2</sub>O, RT;  
then  $Me_3SiCHN_2$   
10) DMDO (1.1 equiv), acetone,

D

Please give a detailed mechanism of step 6.  
Illustrate the stereochemical outcome by a 3D drawing of the transitionstate.



DMDO = dimethyldioxirane

